

# Analysis of the exclusive semileptonic decay $\Lambda_b \rightarrow \Lambda_c + \ell \bar{\nu}_\ell$ within a light-front constituent quark model<sup>a</sup>

F. Cardarelli and S. Simula

Istituto Nazionale di Fisica Nucleare, Sezione Roma III  
Via della Vasca Navale 84, I-00146, Roma, Italy

## Abstract

The exclusive semileptonic decay  $\Lambda_b \rightarrow \Lambda_c + \ell \bar{\nu}_\ell$  is investigated including both radiative and first-order power corrections in the inverse heavy-quark mass, while the Isgur-Wise function is calculated within a light-front constituent quark model. It turns out that the dependence on the model parameters can be effectively constrained using recent lattice *QCD* results at low values of the recoil. Our final predictions for the exclusive semileptonic branching ratio, the longitudinal and transverse asymmetries, and the longitudinal to transverse decay ratio are:  $Br(\Lambda_b \rightarrow \Lambda_c \ell \bar{\nu}_\ell) = (6.3 \pm 1.6) \% |V_{bc}/0.040|^2 \tau(\Lambda_b)/(1.24 \text{ ps})$ ,  $a_L = -0.945 \pm 0.014$ ,  $a_T = -0.62 \pm 0.09$  and  $R_{L/T} = 1.57 \pm 0.15$ , respectively. Moreover, the theoretical uncertainties both on  $a_L$  and the (partially integrated)  $R_{L/T}$  are found to be quite small and, therefore, the experimental determination of these quantities is a very interesting tool for testing the Standard Model and for investigating possible New Physics. In this respect the sensitivity to extract unique information both on the strength and phase of possible hadronic right-handed currents is illustrated.

---

<sup>a</sup>In the Proceedings of the International Conference PANIC '99, Uppsala (Sweden), June 1999, to appear in Nuclear Physics A.



# 1 INTRODUCTION

The investigation of the exclusive semileptonic decays of heavy hadrons, driven by the elementary process  $b \rightarrow c + \ell \bar{\nu}_\ell$ , can provide relevant information on fundamental parameters of the Standard Model ( $SM$ ), like e.g. the quark masses and the weak mixing angle  $V_{bc}$ , as well as on possible extensions of the  $SM$ . In this contribution we present the main results of an analysis of the decay  $\Lambda_b \rightarrow \Lambda_c + \ell \bar{\nu}_\ell$  performed recently in Refs. [1, 2] adopting a relativistic quark model formulated on the light-front, and we illustrate also the possibility to extract relevant information on hadronic right-handed currents from the  $\Lambda_b \rightarrow \Lambda_c + \ell \bar{\nu}_\ell$  process.

In Refs. [1, 2] the Isgur-Wise ( $IW$ ) form factor relevant for the  $\Lambda_b \rightarrow \Lambda_c + \ell \bar{\nu}_\ell$  decay has been calculated in the whole accessible kinematical range adopting a light-front constituent quark model and using various forms of the three-quark wave function. It turns out that the  $IW$  form factor is sensitive to relativistic delocalization effects associated to the light-quark degrees of freedom, leading to a saturation property of the  $IW$  form factor as a function of the canonical baryon size. Moreover, the shape of the  $IW$  function is found to be significantly affected by the baryon structure, being sharply different in case of diquark-like or collinear-type configurations. The comparison with recent lattice  $QCD$  calculations [3] as well as with the (model-dependent) dispersive bounds of Ref. [4] suggests clearly the dominance of collinear-type configurations with respect to diquark-like ones, and allows to put effective *constraints* on the shape of the  $IW$  function in the full recoil range relevant for the  $\Lambda_b \rightarrow \Lambda_c + \ell \bar{\nu}_\ell$  decay (see Ref. [2] for more details).

## 2 ANALYSIS OF THE $\Lambda_b \rightarrow \Lambda_c + \ell \bar{\nu}_\ell$ DECAY WITHIN THE SM

The  $\Lambda_b \rightarrow \Lambda_c + \ell \bar{\nu}_\ell$  decay has been investigated within the framework of the  $SM$  including radiative as well as first-order  $1/m_Q$  corrections to the relevant form factors, as derived in Ref. [5]. Our final predictions for the exclusive semileptonic branching ratio, the longitudinal and transverse asymmetries, and the longitudinal to transverse decay ratio are:  $Br_{SL} = (6.3 \pm 1.4) \% |V_{bc}/0.040|^2 \tau(\Lambda_b)/(1.24 \text{ ps})$ ,  $a_L = -0.94 \pm 0.01$ ,  $a_T = -0.62 \pm 0.08$  and  $R_{L/T} = 1.57 \pm 0.12$ , respectively. Our results for the partially-integrated exclusive semileptonic branching ratio are reported in Fig. 1 and compared with the corresponding results from the lattice  $QCD$  simulations of Ref. [3]. It can be seen that our results are always well within the range of values given by the lattice  $QCD$  simulations and that radiative plus first-order power corrections modify only slightly the results obtained within the Heavy Quark Symmetry ( $HQS$ ). If the integration over the recoil is limited to  $\omega = 1.2$ , then the resulting uncertainty on the branching ratio reduces significantly to  $\simeq 12\%$ , though at the price of reducing the number of the events by a factor  $\simeq 0.44$  (see Table 2 of Ref. [2]). This implies the possibility to extract the  $CKM$  matrix elements  $|V_{bc}|$  with a theoretical uncertainty of  $\simeq 6\%$ , which is comparable with present uncertainties obtained from exclusive semileptonic  $B$ -meson decays [7].




Figure 1. Partially-integrated branching ratio  $Br_{\Lambda_b \rightarrow \Lambda_c \ell \bar{\nu}_\ell}(\omega_{max})$  in % versus the upper limit of integration over the recoil,  $\omega_{max}$ , calculated at  $|V_{bc}| = 0.040$  and  $\tau(\Lambda_b) = 1.24$  ps [6]. The solid and dashed lines correspond to our and lattice  $QCD$  results [3], respectively. The lower and upper solid lines are the results corresponding to the lower and upper limits of the  $IW$  function derived in Ref. [2], including radiative plus first-order  $1/m_Q$  corrections. The dotted lines are the  $HQS$  results. The vertical dot-dashed line indicates the physical threshold  $\omega_{th} \simeq 1.44$ . (After Ref. [2]).

In Ref. [2] various partially-integrated asymmetries have been investigated, showing that in comparing with (future) data the precise  $\omega$ -range of the experiments has to be taken into account. It turns out that radiative plus first-order  $1/m_Q$  corrections are relevant for the transverse asymmetry  $a_T$  and for the longitudinal  $\Lambda_c$  polarization  $P_L$ , whereas both  $a_L$  and  $R_{L/T}$  are only marginally affected by such  $HQS$  corrections, as it is clearly shown in Fig. 2. Moreover, the model dependence on the various asymmetries is generally quite limited and, in particular, our uncertainty on  $R_{L/T}$ , which is always much less than the one presently achievable by lattice  $QCD$  calculations, reduces to  $\simeq 1\%$  when the integration over the recoil is limited to  $\omega = 1.2$ .

### 3 EFFECTS FROM POSSIBLE RIGHT-HANDED CURRENTS

The small uncertainties found for the longitudinal asymmetry  $a_L$  and the longitudinal to transverse decay ratio  $R_{L/T}$  make the experimental determination of these quantities a very interesting tool for testing the  $SM$  and investigating possible New Physics ( $NP$ ). We now illustrate the sensitivity of  $a_L$  and  $R_{L/T}$  to the introduction of possible hadronic right-handed currents by considering the presence in the effective weak Hamiltonian of both left-handed ( $LH$ ) and right-handed ( $RH$ ) operators, viz.  $\mathcal{H}_{eff}(b \rightarrow c \ell \bar{\nu}_\ell) = c_L O_L(b \rightarrow c \ell \bar{\nu}_\ell) + c_R O_R(b \rightarrow c \ell \bar{\nu}_\ell)$ , where  $O_{L(R)}(b \rightarrow c \ell \bar{\nu}_\ell) \equiv (\bar{c} \gamma_\mu (1 \mp \gamma_5) b)(\ell \gamma^\mu (1 - \gamma_5) \nu_\ell)$ , with the values of the coefficients  $c_L$  and  $c_R$  depending on the specific  $NP$  model. In general  $c_L \neq c_L^{(SM)}$ , but in what follows we consider for simplicity  $c_L = c_L^{(SM)}$ , and we introduce two parameters,  $\rho$  and  $\eta$ , defined as  $\rho \equiv |c_R|/|c_L|$  and  $\eta \equiv -\text{Re}(c_L c_R^*)/(|c_L|^2 + |c_R|^2)$ , which are clearly connected to the relative strength and phase of  $RH$  currents with respect to  $LH$  ones. On one hand side we have




Figure 2. Partially-integrated longitudinal asymmetry  $a_L$  (a) and longitudinal to transverse decay ratio  $R_{L/T}$  (d) versus the upper limit of integration over the recoil,  $\omega_{max}$ , for the decay process  $\Lambda_b \rightarrow \Lambda_c + \ell \bar{\nu}_\ell$ . The dotted and solid lines correspond to the  $HQS$  results and to those obtained including radiative plus first-order  $1/m_Q$  corrections, respectively. The meaning of the lower and upper lines, as well as of the vertical dot-dashed line, is as in Fig. 1. In (b) the dashed lines correspond to the lattice  $QCD$  results of Ref. [3]. (Adapted from Ref. [2]).

found that the branching ratio  $Br_{SL}$  is sharply sensitive to the values of  $\rho$  and  $\eta$ ; however, the  $SM$  predictions for  $Br_{SL}$  (see Fig. 1) can be obtained also with different pairs of values of  $(\rho, \eta)$ . On the other hand side our main results for  $a_L$  and  $R_{L/T}$  are summarized in Fig. 3. It can be clearly seen that both  $a_L$  and  $R_{L/T}$  are significantly sensitive to the presence of  $RH$  currents, but their determination allows the extraction of information both on the strength and phase of  $RH$  currents.

## 4 CONCLUSIONS

The exclusive semileptonic decay  $\Lambda_b \rightarrow \Lambda_c + \ell \bar{\nu}_\ell$  has been investigated including both radiative and first-order power corrections in the inverse heavy-quark mass, while the Isgur-Wise function has been calculated within a light-front constituent quark model. The dependence on the model parameters has been effectively constrained using recent lattice  $QCD$  results at low values of the recoil. Our predictions for the exclusive semileptonic branching ratio, the longitudinal and transverse asymmetries, and the longitudinal to transverse decay ratio are:  $Br(\Lambda_b \rightarrow \Lambda_c \ell \bar{\nu}_\ell) = (6.3 \pm 1.6) \% |V_{bc}/0.040|^2 \tau(\Lambda_b)/(1.24 \text{ ps})$ ,  $a_L = -0.945 \pm 0.014$ ,  $a_T = -0.62 \pm 0.09$  and  $R_{L/T} = 1.57 \pm 0.15$ , respectively. Moreover, the theoretical uncertainties both on the longitudinal asymmetry and the (partially integrated) longitudinal to transverse decay ratio are found to be quite small and, therefore, the experimental determination of these quantities is a very interesting tool for testing the Standard Model and for investigating possible New Physics. In this respect the possibility to extract unique information both on the strength and phase of hadronic right-handed currents has been illustrated.




Figure 3. The longitudinal asymmetry  $a_L$  versus the partially-integrated longitudinal to transverse ratio  $\bar{R}_{L/T} \equiv R_{L/T}(\omega_{max} = 1.2)$  for the decay process  $\Lambda_b \rightarrow \Lambda_c + \ell \bar{\nu}_\ell$ . The full dot corresponds to the results calculated within the *SM* framework, while the various lines are obtained for fixed values of  $\rho$  and varying  $\eta$  in its allowed range (see text). The dotted, dashed, dot-dashed, long-dashed, triple-dotted-dashed and solid lines correspond to  $\rho = 0.05, 0.10, 0.15, 0.20, 0.25$  and  $0.30$ , respectively.

## References

- [1] F. Cardarelli and S. Simula: Phys. Lett. B421 (1998) 295.
- [2] F. Cardarelli and S. Simula: preprint INFN-RM3 98/5, e-print archive hep-ph/9810414, to appear in Phys. Rev. D.
- [3] *UKQCD* collaboration, K.C. Bowler et al.: Phys. Rev. D57 (1998) 6948.
- [4] D. Chakraverty et al.: Int. J. Mod. Phys. A14 (1999) 2385.
- [5] M. Neubert: Phys. Rept. 245 (1994) 259.
- [6] Particle Data Group, C. Caso et al., Eur. Phys. J. C3 (1998) 1.
- [7] For an updated analysis of the theoretical uncertainties affecting the extraction of  $V_{bc}$  see I.I. Bigi et al.: Annu. Rev. Nucl. Part. Sci. 47 (1997) 591.